

Preemie or Low Birth Weight after abortion

There have been **40** Statistically Significant Abortion-Preemie and Abortion-LBW studies **for the last 10 years**
(1 Jan. 2005 – 31 December 2014)

1* Chaman R, Amiri M, Ajami M-E, Sadeghian A, Khosravi A.
Low Birth Weight and Its Related Risk Factors in Northeast Iran.
Iran J Pediatrics December 2013;23(6):701-704
[URL:<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4025130/pdf/IJPD-23-701.pdf>]

2 Makhlof MA, Clifton RG, Roberts JM, Myatt L, et al.
Adverse Pregnancy Outcomes among Women with Prior Spontaneous
or Induced Abortions. American J Perinatology 2014
[Abstract URL:
<https://www.thieme-connect.com/ejournals/abstract/10.1055/s-0033-1358771>]

3 McCarthy F, Khashan AS, North RA, Rahma M, Walker JJ, et al.
Pregnancy loss managed by cervical dilation and curettage increases the risk of
spontaneous preterm birth. Human Reproduction Advanced
Access 19 September 2013 Vol. 0 No. 0 pp. 1-10 abstract URL:
<http://www.ncbi.nlm.nih.gov/pubmed/24052504>

4 Oliver-Williams C, Fleming M, Monteath K., Wood AM, Smith GCS.
Changes in Association between Previous Therapeutic Abortion and
Preterm Birth in Scotland, 1980 to 2008: A Historical Cohort Study.
PLOS July 2013;10(7):1-11
<http://www.plosmedicine.org/article/fetchObject.action?uri=info%3A%2F10.1371%2Fjournal.pmed.1001481&representation=PDF>
[Although the 2013 'Oliver-Williams' study found significantly elevated
risk of preterm birth, for the 21st century this study did NOT find
elevated preterm birth risk for Scottish women with prior induced
abortions]

5 Scholten BL, Page-Christiaens CML, Franx A, Hukkelhoven
CWPM, Koster MPH. The influence of pregnancy termination on the
outcome of subsequent pregnancies: a retrospective cohort study.
BMJ OPEN 2013;3:e002803 doi:10.1136/bmjopen-2013-002803
<http://bmjopen.bmj.com/content/3/5/e002803.full.pdf+html>

6 Raisanen S, Gissler M, Saari J, Kramer M, Heinon S.
Contribution of Risk Factors to Extremely, Very and Moderately
Preterm Term Births- Register-Based Analysis of 1,390,742 Singleton

Births. PLOS ONE April 2013;8(4):1-7

<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0060660>

7 Watson LF, Rayner J-A, Forster D. Identifying risk factors for very preterm birth: A reference for clinicians. *Midwifery* May 2013; 29(5):434-439

8 Ghislain Hardy, Alice Benjamin, Haim A. Abenhaim. Effects of Induced Abortions on Early Preterm Births and Adverse Perinatal Outcomes. *Journal Obstetrics Gynaecology Canada* 2013;35(2):138-143 [URL: http://www.jogc.com/abstracts/full/201302_Obstetrics_5.pdf]

9* Heaman M, Kingston D, Chalmers B, Sauve R, Lee L, Young D. Risk Factors for Preterm Birth and Small-for-gestational-age among Canadian Women. *Paediatric Perinatal Epidemiology* 2013;27:54-61

10 Klemetti R, Gissler M, Niinimäki M, Hemminki E. Birth outcomes after induced abortion: a nationwide register-based study of first births in Finland. *Human Reproduction* 2012 August 29 [Epub ahead of print] <http://www.ncbi.nlm.nih.gov/pubmed/22933527>

11 Bhattacharya S, Lowit A, Bhattacharya S, Raja EA, Lee AM, Mahmood T, Templeton A. Reproductive outcomes following induced abortion; a national register-based cohort study in Scotland. *BMJ OPEN* 2012;2:e000911 [doi:10.1136/bmjopen-2012-000911 ; URL: <http://bmjopen.bmj.com/content/2/4/e000911.full.pdf>]

12 Renzo GCD, Giardina I, Rosati A, Clerici G, Torricelli M, et al. Maternal risk factors for preterm birth: a country based population analysis. *European J Obstetrics Gynecology and Reproductive Biology*. December 2011;159(2):342-346 URL: <http://www.sciencedirect.com/science/article/pii/S0301211511005392>

13 Liao H, Wei Q, Duan L, Ge J, Zhou Y, Zeng W. Repeated medical abortions and the risk of preterm birth in subsequent pregnancies. *Arch Gynecol Obstet* 2011;289:579-586 URL: <http://www.springerlink.com/content/d5mt821806512570/>

14 Furquim MA, Alencar GP, Schoeps D, Novaes HMD, Campbell O, et al. Survival and risk factors nor neonatal mortality in a cohort of very low birth weight infants in the southern region of San Paulo city, Brazil. *Cadernos de Saude Publica* 2011;27(6)

15 Junli G, Weiyue Z. Influence of artificial abortion on preterm labor and risk in subsequent pregnancy. Chinese J Obstet Gynaecol Pediatrics [Electronic] 2010;447-451

16 Ammar M Alfidhli, Ali M Hajia, Farida AK Mohammed, Hamdiya A Alfidhli, Medhat K El-Shazly. Incidence and Potential Risk Factors of Low Birth Weight Among Full Term Deliveries 2010;46(2):157-164 [Study Population: Kuwaiti women]

17+ Watson LF, Rayner J-A, King J, Jolley D, Forster D, Lumley J. Modelling sequence of prior pregnancies on subsequent risk of very preterm birth. Paediatric and Perinatal Epidemiology 2010;24:416-423

18+ Watson LF, Rayner J-A, King J, Jolley D, Forster D, Lumley J. Modelling prior reproductive history to improve predication of risk for very preterm birth. Paediatric Perinatal Epidemiology 2010;24:402-415

19 Yuan W, Duffner AM, Chen L, Hunt LP, Sellers SM, Bernal AL. Analysis of preterm deliveries below 35 weeks' gestation in a tertiary hospital in the UK. A case-control survey.. BMC Research Notes 2010;3:119

2000-2009

20 Voigt M, Henrich W, Zygmunt M, Friese K, Straube S, Briese V. Is induced abortion a risk factor in subsequent pregnancy? Journal Perinatal Medicine 2009;37:144-149 [Study Population: German women]

21 Freak-Poli R, Chan A, Gaeme J, Street J. Previous abortion and risk of preterm birth: a population study. J Maternal-Fetal Med Jan. 2009; 22(1):1-7 [Study Population: Australian women]

22 Hldre K, Rahu K, Karro H, Rahu M. Previous history of surgically induced abortion and complications of the third stage of labour in subsequent normal vaginal deliveries. J Maternal-Fetal Neonatal Medicine 2008;21(12): 884-888 [Study Population: Estonian women]

23 Visintine J, Berghella V, Henning D, Baxter J. Cervical length for prediction of preterm birth in women with multiple prior induced abortions. Ultrasound Obstetrics Gynecology 2008;31(2):198-200 [Study Population: U.S. women]

24* Sareer Badshah, Linda Mason, Kenneth McKelvie, Roger Payne, Paulo JG Lisboa. Risk factors for low birthweight in the public-hospitals at Peshawar NWFP-Pakistan 2008;8:197-206 [Study Population: Pakistani women]

25 Voigt M, Olbertz D, Fusch C, Krafczyk D, Briese V, Schneider KT. The influence of previous pregnancy terminations, miscarriages, and still-birth on the incidence of babies with low birth weight and premature births as well as somatic classification of newborns.

Z Geburtshilfe Neonatol 2008;212:5-12 [Study Population:
German women]

26 Reime B, Schuecking BA, Wenzlaff P. Reproductive Outcomes in
Adolescents Who Had a Previous Birth or an Induced Abortion Compared
to Adolescents' First Pregnancies. BMC Pregnancy and Childbirth
2008;8:4 [Study Population: German women]

27* Brown TS, Adera T, Masho SW. Previous abortion and the risk of
low birth weight and preterm births. J Epidemiol Commun Health
2008;62:16-22 [Study Population: U.S. women]

28 Curry AE, Vogel I, Drews C, Schendel D, Skogstrand K, et al.
Mid-pregnancy maternal plasma levels of interleukin 2, 6, and 12, tumor
necrosis factor-alpha, interferon-gamma, and granulocyte-macrophage
colony-stimulating factor and spontaneous preterm delivery. Acta
Obstet et Gynecologica 2007;86:1103-1110 [Study Population:
Danish women]

29 Chung-Chin Lo, Jenn-Jeih Hsu, Ching-Chang Hsieh, T'sang-T-sang
Hsieh, Tai-Ho Hung. Risk Factors For Spontaneous Preterm Delivery
Before 34 Weeks of Gestation Among Taiwanese Women. Taiwan J
Obstet Gynecol 2007;46(4):389-394 [Study Population: Taiwanese
women]

29 Jackson JE, Grobman WA, Haney E, Casele H. Mid-trimester
dilation and evacuation with laminaria does not increase the
risk for severe subsequent pregnancy complications. Intl J
Gynecol Obstet 2007;96:12-15 [Study Population: U.S. women]

30 Briunsmas F, Lumley J, Tan J, Quinn M. Precancerous changes
in the cervix and risk of subsequent preterm birth. BJOG
Jan. 2007;114(1):70-80 { Study Population: Australian women]

31 Losa SM, Gonzalez E, Gonzalez G. Risk Factors for Preterm
Birth. Prog Obstet Ginecol 2006;49(2):57-65

32 Teramoto S, Soeda A, Hayashi Y, Urashima M. Physical and
socioeconomic predictors of birth weight in Japan. Pediatrics
International 2006;48(3):274-277

33 Smith GCS, Shah I, White IR, Pell JP, Crossley JA, Dobbie R.
Maternal and biochemical predictors of spontaneous preterm birth
among nulliparous women: a systematic analysis in relation to
degree of prematurity. Intl J Epidem 2006;35(5):1169-1177
[Study Population: Scottish women]

35 K.K. Roy, Jinee Baruah, Sunesh Kumar, Neena Malhotra,
A.K. Deorari, J.B. Sharma. Maternal Antenatal Profile and
Immediate Neonatal Outcome in VLBW and ELBW Babies.

Indian Journal of Pediatrics 2006;73(8):669-673
[Study Population: Indian women in India]

36 Samin A, Al-Dabbagh, Wafa Y Al-Tae. Risk factors for preterm birth in Iraq: a case-control study. Pregnancy and Childbirth. BMC 2006;6:13 [Study Population: women in Iraq]

37 Poikkens P, Unkila-Kallio L, Vilska S, Repokari L. et al. Impact of Infertility Characteristics and treatment modalities on singleton pregnancies after assisted reproduction. Reproductive Biomed July 2006;13(1):135-144 [Study Population: Finnish women]

38 Etuk SJ, Etuk IS, Oyo-Ita AE. Factors Influencing the Incidence of Pre-term Birth in Calabar, Nigeria. Nigerian J Physiological Sciences 2005;20(1-2):63-68 [Study Population: Nigerian women]

39 Stang P, Hammond AO, Bauman P. Induced Abortion Increases the Risk of Very Preterm Delivery; Results from a Large Perinatal Database. Fertility Sterility Sept 2005;S159 [Study Population: German women]

40 Conde-Agudelo A, Belizan JM, Breman R, Brockman SC, Rosas-Bermudez. Effect of the interpregnancy interval after an abortion on maternal and perinatal health in Latin America. Int J Gynaecol & Obstet 2005;89 (Sup. 1):S34-S40. [Study Population: from Latin American countries]

.....
* - studies in which 'abortion' means spontaneous abortion (i.e. miscarriage) or induced abortion

Compiled by:
info@abortioninformation.eu

Data from:
Brent Rooney <fullterm40@gmail.com>